PROFESSOR JAYASHANKAR TELANGANA STATE AGRICULTURAL UNIVERSITY

Agro Climate Research Centre (ACRC),
Agricultural Research Institute, Rajendranagar, Hyderabad-30.

Bulletin No. XXXIII/39/2022

WEATHER BASED AGROMET ADVISORIES FOR TELANGANA STATE

FROM DATE: 18.05.2022 (Wednesday) TO 22.05.2022 (Sunday)

Grameen Krishi Mausam Seva (GKMS Project)

INDIA METEOROLOGICAL DEPARTMENT, NEW DELHI.

Weather details of past 3 days:

During the last three days light to moderate rains occurred at isolated places over different parts of the State. The maximum and minimum temperatures ranged between 35-44^{0}C and 21-31^{0}C respectively.

Forecasted weather for the next 5 days:

As per the forecast received from Meteorological Centre, Hyderabad, light to moderate rains may occur at isolated places over different parts of the State during next five days. The maximum and minimum temperatures are likely to range between 37-42^{0}C and 24-30^{0}C respectively.

Warning:

Thunderstorm accompanied with lightening and gusty winds (30-40 kmph) very likely to occur at isolated places in Telangana during the next 24 hours.

Weather based Agromet Advisories

- Southwest Monsoon has advanced into some parts of South Bay of Bengal, most part of Andaman Nicobar Islands and Andaman Sea as on 16th May, 2022.
- In view of the thunderstorm activity and rainfall forecast during the next five days. Farmers are advised to
  - Cover the already harvested crop produce with tarpaulin to avoid post harvest losses.
- By taking the advantage of rains farmers are advised to take up summer ploughing.
- Dig the pits for planting new orchards.

Rice

- Harvesting, Threshing, Storage of grain etc operations in paddy may be carried out by observing the weathering conditions.

Vegetables

- If incidence of sucking pests is noticed. To control, spray Dimethoate @ 2 ml or Acephate @ 1.5 g per litre of water.
- Prevailing weather conditions are congenial for Leaf spot in vegetable crops. To control, spray Carbendazim @ 1 g or Propiconazole @1 ml per liter of water twice in 15 days interval.
- Incidence of Shoot and Fruit Borer is noticed in Brinjal. To control,
  - Installation of pheromone traps to monitor the fruit and shoot borer

- Clip the infested shoots and destroy
- Spray Profenophos @ 2 ml or Emamectin Benzoate @ 0.4g per liter of water

- Incidence of Chilli flower thrips is noticed. To control, spay Fipronil @ 2 ml per liter of water.

Mango

- Prevailing weather conditions are congenial for the incidence of fruit fly in mango. To trap the flies in the container, hang the plastic containers with mixture of methyl eugenol @ 2 ml + Carbofuran 3G @ 3g per litre of water in the orchard.
- Incidence of Hoppers, Mealy Bugs and Scale Insects are noticed in Mango due to recent rains. To control, spray Imidachloprid @ 0.3 ml + Neem oil @ 2.5 ml per liter of water.

Poultry

- Prevailing weather conditions are congenial for the occurrence of Ranikhet disease in poultry. To prevent the diseases, vaccinate the birds.
- Protect the poultry birds by arranging fans and foggers in sheds, cover the sheds with paddy straw and arrange sprinklers. Feed with wet mash to increase the feed intake and provide cool water for drinking.

Livestock

- Prevailing weather conditions are congenial for the occurrence of
  - Ranikhet disease in poultry, ET, sheep pox and PPR in Sheep
  - Hemorrhagic Septicemia, Foot and mouth disease in cattle and Buffaloes. To prevent the diseases, vaccinate the animals

Senior Scientist (Agro) & Head
Agro Climate Research Centre (ACRC),
PJTSAU, Rajendranagar, Hyd-30.

ప్రొఫెసర్ జయశంకర్ తెలంగాణ రాష్ట్ర వ్యవసాయ విశ్వవిద్యాలయం
వ్యవసాయ వాతావరణ పరిశోధన కేంద్రం
వ్యవసాయ పరిశోధన సంస్థ, రాజేంద్రనగర్, హైదరాబాద్.

బులెటిన్ నెం. XXXIII/39/2022

రాష్ట్ర రైతాంగానికి వాతావరణ ఆధారిత వ్యవసాయ సలహాలు

తేది: 18.05.2022 (బుధవారం) నుండి 22.05.2022 (ఆదివారం) వరకు

గ్రామీణ కృషి మౌసమ్ సేవ పథకం (GKMS Project)
భారత వాతావరణ శాఖ, న్యూఢిల్లీ.

గత మూడు రోజుల వాతావరణం:

గడిచిన మూడు రోజులలో రాష్ట్రంలో అక్కడక్కడ తేలికపాటి నుండి ఓ మోస్తరు వర్షాలు కురిసాయి. పగటి ఉష్ణోగ్రతలు 35-44 డిగ్రీల సెల్సియస్ మరియు రాత్రి ఉష్ణోగ్రతలు 21-31 డిగ్రీల సెల్సియస్ గా నమోదయ్యాయి.

రాబోవు ఐదు రోజుల వాతావరణం:

హైదరాబాద్ వాతావరణ కేంద్రం వారు అందించిన సమాచారం ప్రకారం రాబోవు ఐదు రోజులలో రాష్ట్రంలో అక్కడక్కడ తేలికపాటి నుండి ఓ మోస్తరు వర్షాలు కురిసే సూచనలున్నాయి. పగటి ఉష్ణోగ్రతలు 37-42 డిగ్రీల సెల్సియస్ మధ్య మరియు రాత్రి ఉష్ణోగ్రతలు 24-30 డిగ్రీల సెల్సియస్ మధ్య నమోదుకావచ్చు.

హెచ్చరిక:

రాబోవు 24 గంటలలో తెలంగాణలో అక్కడక్కడ ఉరుములతో కూడిన మెరుపులు మరియు ఈదురు గాలులు (30-40 కి. మీ /గంటకు) వీచే అవకాశం ఉంది.

వాతావరణ ఆధారిత వ్యవసాయ సలహాలు

- మే 16, 2022 రోజు వరకు నైరుతి ఋతుపవనాలు దక్షిణ బంగాళాఖాతంలో కొన్ని ప్రాంతాలలో మరియు అండమాన్ నికోబార్ అంతటా వ్యాపించాయి.
- రాబోవు ఐదు రోజులలో ఉరుములతో కూడిన వర్ష సూచనలు ఉన్నందున, రైతులు
  - కోసిన పంటలను తార్పలిన్ తో కప్పి భద్రపర్చుకోవాలి.
- గత కొన్ని రోజులుగా రాష్ట్రంలో అక్కడక్కడ కురుస్తున్న వర్షాలను ఉపయోగించుకొని వేసవి దుక్కులు చేసుకోవాలి.
- క్రొత్త పండ్ల తోటలు పెట్టడానికి గుంతలు త్రవ్వుకోవాలి.

వరి

- వాతావరణ పరిస్థితులను గమనిస్తూ పంటకోత, నూర్పిడి, ధాన్యము నిల్వ చేయుట మొదలగు పనులను చేసుకోవాలి.

కూరగాయలు

- కూరగాయ పంటలలో రసం పీల్చే పురుగులు గమనించినట్లైతే, నివారణకు 2 మి.లీ. డైమిథోయేట్ లేదా 1.5 గ్రా. ఎసిఫేట్ లీటరు నీటికి కలిపి పిచికారి చేయాలి.
- ప్రస్తుత వాతావరణ పరిస్థితులు కూరగాయల పంటలలో ఆకుమచ్చ తెగులు సోకుటకు అనుకూలము. తెగులు నివారణకు 1 గ్రా. కార్బెండజిమ్ లేదా 1 మి.లీ. ప్రోపికోనజోల్ మందును లీటరు నీటిలో కలిపి 15 రోజుల వ్యవధిలో రెండు సార్లు పిచికారి చేయాలి.
- వంగలో కొమ్మ మరియు కాయతొలుచు పురుగు గమనించడమైనది. నివారణకు,
  - లింగాకర్షక బుట్టలను అమర్చి పురుగు యొక్క ఉధృతిని పర్యవేక్షించుకోవాలి.
  - పురుగు సోకిన కొమ్మలను త్రుంచి నాశనం చేయాలి.

- 2 మి.లీ. ప్రొఫెనోఫాస్ లేదా 0.4 గ్రా. ఇమమెక్టిన్ బెంజోయేట్ మందును ఒక లీటరు నీటిలో కలిపి పిచికారి చేయాలి.

- మిరపలో తామరపురుగు పూతను ఆశించడమైనది. నివారణకు, 2 మి.లీ. ఫిప్రోనిల్ మందును లీటరు నీటికి కలిపి పిచికారి చేయాలి.

మామిడి

- ప్రస్తుత వాతావరణ పరిస్థితులలో మామిడిలో పండు ఈగ గమనించడమైనది. నివారణకు, ప్లాస్టిక్ పళ్ళెంలో 2 మి.లీ. మిథైల్ యూజినాల్ మరియు 3 గ్రా. కార్బోఫ్యూరాన్ 3 జి గుళికలను లీటరు నీటికి కలిపి తోటలో వేలాడదీసినచో పురుగులు మందు నీటిలో పడి చనిపోతాయి.
- మామిడిలో తేనె మంచు పురుగు, పిండి నల్లి మరియు పొలుసు పురుగుల ఉధృతి గమనించడమైనది. నివారణకు, 0.3 మి.లీ. ఇమిడాక్లోప్రిడ్ మందును + 2.5 మి.లీ వేపనూనెను లీటరు నీటికి కలిపి పిచికారి చేయాలి.

కోళ్ళు

- ప్రస్తుత వాతావరణ పరిస్థితులు కోళ్ళలో కొక్కెర తెగులు సోకుటకు అనుకూలం. నివారణకు టీకాలు వేయించాలి.
- కోళ్ళు వేడి వాతావరణాన్ని తట్టుకొనుటకు షెడ్లల్లో ఫ్యాన్లను మరియు ఫాగార్స్ ను అమర్చి షెడ్లపై వరిగడ్డితో కప్పి స్ప్రింక్లర్లను అమర్చాలి. కోళ్ళు ఎక్కువ మోతాదులో తినుటకు అనుగుణంగా మెత్తటి దానాను పెట్టి త్రాగుటకు చల్లని నీటిని అందుబాటులో ఉంచాలి.

పశుపోషణ

- ప్రస్తుత వాతావరణ పరిస్థితులు కోళ్ళలో కొక్కెర తెగులు సోకుటకు అనుకూలం వీటి నివారణకు, టీకాలు వేయించాలి.
- గొర్రెలలో చిటుకు, మశూచి వ్యాధి మరియు PPR
- ఆవులు మరియు గేదేలలో గొంతువాపు వ్యాధి, గొంతు మరియు గాలికుంటు వ్యాధి, సోకుటకు అనుకూలం వీటి నివారణకు, టీకాలు వేయించాలి.

సీనియర్ శాస్త్రవేత్త (అగ్రానమి)& అధిపతి

ప్రకటన సీనియర్ శాస్త్రవేత్త, వ్యవసాయ వాతావరణ పరిశోధన కేంద్రం వ్యవసాయ పరిశోధన సంస్థ రాజేంద్రనగర్
ఫోన్. నెం. 9989625231