

ప్రాఫెసర్ జయశంకర్ తెలంగాణ రాష్ట్ర వ్యవసాయ విశ్వవిద్యాలయం

వ్యవసాయ వాతావరణ పరిశోధన కేంద్రం

వ్యవసాయ పరిశోధన సంస్థ, రాజేంద్రనగర్, హైదరాబాద్.



బులెటిన్ నెం. XXXIII/47/2022

రాష్ట్ర రైతాంగానికి

వాతావరణ ఆధారిత వ్యవసాయ సలహాలు

తేది: 15.06.2022 (బుధవారం) నుండి 19.06.2022 (ఆదివారం) వరకు



గ్రామీణ కృషి మౌసమ్ సేవ పథకం (GKMS Project)

భారత వాతావరణ శాఖ, న్యూఢిల్లీ.

## గత మూడు రోజుల వాతావరణం:

గడిచిన మూడు రోజులలో రాష్ట్రంలో చాలా చోట్ల తేలికపాటి నుండి ఓ మోస్తరు వర్షాలు కురిసాయి. పగటి ఉష్ణోగ్రతలు 35-42 డిగ్రీల సెల్సియస్ మరియు రాత్రి ఉష్ణోగ్రతలు 20-30 డిగ్రీల సెల్సియస్ గా నమోదయ్యాయి.

## రాబోవు ఐదు రోజుల వాతావరణం:

హైదరాబాద్ వాతావరణ కేంద్రం వారు అందించిన సమాచారం ప్రకారం రాబోవు ఐదు రోజులలో రాష్ట్రంలో చాలా ప్రాంతాలలో తేలికపాటి నుండి ఓ మోస్తరు వర్షాలు కురిసే సూచనలున్నాయి. పగటి ఉష్ణోగ్రతలు 32-40 డిగ్రీల సెల్సియస్ మధ్య మరియు రాత్రి ఉష్ణోగ్రతలు 21-27 డిగ్రీల సెల్సియస్ మధ్య నమోదుకావచ్చు.

## వాతావరణ హెచ్చరిక:

- రాబోవు రెండు రోజులలో రాష్ట్రంలో అక్కడక్కడ భారీ వర్షాలు కురిసే సూచనలున్నాయి.
- రాబోవు ఐదు రోజులలో రాష్ట్రంలో అక్కడక్కడ ఉరుములు మెరుపులతో కూడిన వర్షాలు కురిసే సూచనలున్నాయి.

## వాతావరణ ఆధారిత వ్యవసాయ సలహాలు

- నైరుతి ఋతుపవనాలు రాష్ట్రంలోనికి ప్రవేశించిన తరువాత తేలిక పాటి నేలల్లో 50 నుంచి 60 మి.మీ. మరియు బరువు నేలలో 60 నుంచి 75 మి.మీ వర్షపాతం నమోదైన తరువాత లేదా నేల 15-20 సెం.మీ. లోతు తడిసిన తరువాతనే రైతులు వర్షాధార పంటలను విత్తుకోవాలి.
- సజ్జ మరియు కొర్రలను పంటలను 31 జూలై వరకు విత్తుకోవచ్చును.
- ఆముదం పంటను 31 జూలై వరకు విత్తుకోవచ్చును. కిలో విత్తనానికి 3 గ్రా. థైరమ్ లేదా కాప్టాన్ కలిపి విత్తన శుద్ధి చేయడం వలన మొలక కుళ్ళు తెగుళ్ళు అరికట్టవచ్చు.
- సాయాచిక్కుడు పంటను తేలికపాటి నేలలో సాగు చేయరాదు. ఈ పంటను జూలై 10 తేదీ వరకు విత్తుకోవాలి. కిలో విత్తనానికి 1.5 మి.లీ. ఇమిడాక్లోప్రిడ్ 48%తో మరియు 1.0 గ్రా. కార్బండజిమ్ తో విత్తనశుద్ధి చేసి విత్తుకొంటే తొలిదశలో ఆశించు పురుగులు, తెగుళ్ళు నుండి కావలసిన రక్షణ లభిస్తుంది.
- ప్రత్తి పంటను జూలై 20వ తేదీ వరకు విత్తుకోవచ్చు.

## వరి

- నీరు సమృద్ధిగా ఉన్న ప్రాంతాల్లో రైతులు దీర్ఘకాలిక వరి రకాల నారుమళ్ళు పోసుకోవడానికి ఇది అనువైన సమయం.
- వరిలో 3 గ్రా. కార్బండజిమ్ కిలో విత్తనానికి కలిపి విత్తనశుద్ధికి చేసుకోవాలి.

- వరి సాగు చేసే పొలాల్లో, తొలకరి వర్షాలను ఉపయోగించుకొని జనుము లేదా జీలుగను పచ్చి రొట్ట పైరుగా, ఆలస్యంగా నీరు విడుదలయ్యే ప్రాంతాలలో పెసరను పైరుగా లేనిచో పచ్చి రొట్టగా విత్తుకోవాలి.
- రైతులు తెలంగాణ సాన (RNR 15048) వరి విత్తనాన్ని జూన్ నెలలో నారుమడి పోసుకోకూడదు.

### మొక్కజొన్న

- వర్షాధారంగా పండించే మొక్కజొన్న పైరును విత్తుటకు జూన్ 15 నుండి జూలై ఆఖరు వరకు అనుకూలమైన సమయం. జూన్ 15 నుండి జూలై 15 వరకు దీర్ఘకాలిక మరియు మధ్యకాలిక రకాలు మరియు జూలై 15 నుండి జూలై ఆఖరు వరకు స్వల్పకాలిక రకాలను విత్తుకోవచ్చు.
- మొక్కజొన్నలో కత్తెర పురుగు నివారణకు కిలో విత్తనానికి 6.0 మి.లీ. సాయంబ్రానీలిప్రోల్ + థయోమిథాగ్లామ్ తో విత్తన శుద్ధి చేసుకున్నట్లయితే తొలిదశలో మొక్కలను కత్తెర పురుగు బారినండి కాపాడుకోవచ్చును.

### కూరగాయలు

- కూరగాయ పంటలలో రసం పీల్చే పురుగులు గమనించినట్లైతే, నివారణకు 1.5 గ్రా. ఎసిఫెట్ లీటరు నీటికి కలిపి పిచికారి చేయాలి.
- ప్రస్తుత వాతావరణ పరిస్థితులు కూరగాయల పంటలలో ఆకుమచ్చ తెగులు సోకుటకు అనుకూలము. తెగులు నివారణకు 1 గ్రా. కార్బెండజిమ్ లేదా 1 మి.లీ. ప్రావికోనజోల్ మందును లీటరు నీటిలో కలిపి 15 రోజుల వ్యవధిలో రెండు సార్లు పిచికారి చేయాలి.
- వంగలో కొమ్మ మరియు కాయతొలుచు పురుగు గమనించడమైనది. నివారణకు,
  - లింగాకర్షక బుట్టలను అమర్చి పురుగు యొక్క ఉద్యతని పర్యవేక్షించుకోవాలి.
  - పురుగు సోకిన కొమ్మలను త్రుంచి నాశనం చేయాలి.
  - 2 మి.లీ. ప్రాఫెనోఫాస్ లేదా 0.4 గ్రా. ఇమమెక్టిన్ బెంజోయేట్ మందును ఒక లీటరు నీటిలో కలిపి పిచికారి చేయాలి.
- మిరపలో తామరపురుగు పూతను ఆశించడమైనది. నివారణకు, 2 మి.లీ. ఫిప్రోనిల్ మందును లీటరు నీటికి కలిపి పిచికారి చేయాలి.

### కోళ్ళు

- ప్రస్తుత వాతావరణ పరిస్థితులు కోళ్ళలో కొక్కెర తెగులు సోకుటకు అనుకూలం. నివారణకు టీకాలు వేయించాలి.
- కోళ్ళు వేడి వాతావరణాన్ని తట్టుకొనుటకు షెడల్లో ఫ్యాన్లను మరియు ఫాగర్స్ ను అమర్చి షెడల్లో పరిగడ్డితో కప్పి స్ప్రింక్లర్లను అమర్చాలి. కోళ్ళు ఎక్కువ మోతాదులో తినుటకు అనుగుణంగా మెత్తటి దానాను పెట్టి త్రాగుటకు చల్లని నీటిని అందుబాటులో ఉంచాలి.

## పశుపోషణ

- ప్రస్తుత వాతావరణ పరిస్థితులు కోళ్ళలో కొక్కెర తెగులు సోకుటకు అనుకూలం వీటి నివారణకు, టీకాలు వేయించాలి.
- గొర్రెలలో చిటుకు, మశూచి వ్యాధి మరియు PPR
- ఆవులు మరియు గేదేలలో గొంతువాపు వ్యాధి, గొంతు మరియు గాలికుంటు వ్యాధి, సోకుటకు అనుకూలం వీటి నివారణకు, టీకాలు వేయించాలి.

సీనియర్ శాస్త్రవేత్త (అగ్రానమి)& అధిపతి

---

ప్రకటన: సీనియర్ శాస్త్రవేత్త, వ్యవసాయ వాతావరణ పరిశోధన కేంద్రం, వ్యవసాయ పరిశోధన సంస్థ, రాజేంద్రనగర్ ఫోన్. నెం. 9989625231

**PROFESSOR JAYASHANKAR TELANGANA STATE  
AGRICULTURAL UNIVERSITY**

**Agro Climate Research Centre (ACRC),  
Agricultural Research Institute, Rajendranagar, Hyderabad-30.**



**Bulletin No. XXXIII/47/2022**

**WEATHER BASED AGROMET ADVISORIES  
FOR TELANGANA STATE**

**FROM DATE: 15.06.2022 (Wednesday) TO 19.06.2022 (Sunday)**



**Grameen Krishi Mausam Seva (GKMS Project)**

**INDIA METEOROLOGICAL DEPARTMENT, NEW DELHI.**

### **Weather details of past 3 days:**

During the last three days light to moderate rains occurred at many places over different parts of the State. The maximum and minimum temperatures ranged between 35-42<sup>0</sup>C and 20-30<sup>0</sup>C respectively.

### **Forecasted weather for the next 5 days:**

As per the forecast received from Meteorological Centre, Hyderabad, light to moderate rains very likely to occur at many places over different parts of the State during next five days. The maximum and minimum temperatures are likely to range between 32-40<sup>0</sup>C and 21-27<sup>0</sup>C respectively.

### **Warning:**

- Heavy rains very likely to occur at isolated places over Telangana during the next two days.
- Thunderstorm accompanied with lightening very likely to occur at isolated places in Telangana during the next five days.

### **Weather based Agromet Advisories**

- Sow the rainfed crops only after receiving cumulative rainfall of 50-60 mm in light soils and 60-75 mm in heavy soils or the soil should be wet up to top 15-20 cm after onset of South-West monsoon rains in the State.
- Take up sowing of Bajra and Korra up to 31 July.
- Take up sowing of Castor crop up to 31 July. Seed treatment with Thiram or Captan @ 3g per kg seed to avoid seedling rot in Castor.
- Don't take up sowing of Soybean in light soils. Take up sowing of Soybean up to 10 July. Seed treatment with Imidachloprid 48% @ 1.5 ml and Carbendazim @ 1.0 g to avoid insect pest and diseases during early stages of the crop.
- Take up sowing of Cotton up to 20 July.

### **Rice**

- Farmers are advised to raise rice nurseries of long duration varieties wherever sufficient water is available.
- Paddy seed should be treated with Carbendazim @ 3g per kg seed.
- By taking advantage of the present rains, take-up sowing of Sunhemp or Dhaincha as in-situ green manure crop preceding rice. Sow Greengram as catch crop preceding rice depending on the availability of release of water.
- Don't sow the paddy nurseries using Telangana Sona (RNR 15048) in the month of June.

### **Maize**

- Rainfed maize may be sown from 15 June to 31 July. Long and medium duration hybrids may be sown between 15 June to 15 July and short duration hybrids may be sown between 15 July to 31 July.
- To control Fall Army Worm, seed may be treated with Cyantraniliprole + Thiamethoxam @ 6 ml per kg seed.

### **Vegetables**

- If incidence of sucking pests is noticed. To control, spray Acephate @ 1.5 g per litre of water.
- Prevailing weather conditions are congenial for Leaf spot in vegetable crops. To control, spray Carbendazim @ 1 g or Propiconazole @1 ml per liter of water twice in 15 days interval.
- Incidence of Shoot and Fruit Borer is noticed in Brinjal. To control,
  - Installation of pheromone traps to monitor the fruit and shoot borer
  - Clip the infested shoots and destroy
  - Spray Profenophos @ 2 ml or Emamectin Benzoate @ 0.4g per liter of water
- Incidence of Chilli flower thrips is noticed. To control, spray Fipronil @ 2 ml per liter of water.

### **Poultry**

- Prevailing weather conditions are congenial for the occurrence of Ranikhet disease in poultry. To prevent the diseases, vaccinate the birds.
- Protect the poultry birds by arranging fans and foggers in sheds, cover the sheds with paddy straw and arrange sprinklers. Feed with wet mash to increase the feed intake and provide cool water for drinking.

### **Livestock**

- Prevailing weather conditions are congenial for the occurrence of
  - Ranikhet disease in poultry, ET, sheep pox and PPR in Sheep
  - Hemorrhagic Septicemia, Foot and mouth disease in cattle and Buffaloes.To prevent the diseases, vaccinate the animals

**Senior Scientist (Agro) & Head  
Agro Climate Research Centre (ACRC),  
PJTSAU, Rajendranagar, Hyd-30.**